

The Role of Tannins as Plant Based Nematicide in Root-Gall Nematode Management of Okra (*Abelmoschus esculentus*) for Improved Yield

by Ezechukwu Vincent Ogwudire

Submission date: 26-Jul-2024 10:54PM (UTC+0100)

Submission ID: 2422920972

File name: PHYTOPATHOLOGY_MANUSCRIPT.docx (138.38K)

Word count: 5593

Character count: 31549

The Role of Tannins as Plant Based Nematicide in Root-Gall Nematode Management of Okra (*Abelmoschus esculentus*) for Improved Yield

Ogwudire Ezechukwu Vincent^{1*}, Ojiako Onyemaobi Frank¹, Ewelike Nicholas³ Nwokeji Ephraim¹, Echerobia Ogbuji Christopher¹, Michael Akaninyene Okon², Ejiogu Stephen¹, Kanu Benjamin¹ and Umelo Queen Chidinma⁴

¹Department of Crop Science and Technology, Federal University of Technology, PMB 1520, Owerri, Nigeria.

²Department of Soil Science and Technology, Federal University of Technology, PMB 1520, Owerri, Nigeria

³Department of Microbiology, Federal University of Technology, P.M.B.1526, Owerri, Nigeria.

⁴Department of, University of Agriculture and Environmental Sciences, Umuagwo, Imo State, Nigeria

*For correspondence: vincent.ogwudire@futo.edu.ng

ABSTRACT

In vitro and screen house studies were conducted to investigate the effectiveness of physicinut Tannins and plant parts against root-gall nematode infected okra. Effectiveness of Tannins against nematode juveniles (J2s) mortality was evaluated in the laboratory. Data was collected on percentage mortality at 12, 24 and 72 h incubation periods. Findings revealed the highest nematode mortality of 94 % after 72 h incubation period with root extracts of tannins at 10 ml rate of application. Results obtained under screen house condition showed that Leaf and Seed extracts of Tannins applied at 5 ml/pot and inoculated with 1,200 infective nematode larvae (J2s) significantly ($P=0.05$) decreased root-gall index from 4.00 (severely galled) to 1.00 (rarely galled) while enhancing growth and yield attributes of okra than the untreated control. Maximum effect of treatment rates on root gall index and yield was most impactful at the highest dose of 5ml/pot. Correlation of root-gall index and yield attributes substantiated the effectiveness of Tannins in improving the growth and yield of Okra at reduced galling responses. While leaf and seed Tannins application were most effective in increasing plant heights, leaf areas and okra pod weights by 67 % (4.88 g) over the control, highest 100 dry seed weights and improved mucilaginous property was achieved with root tannins application at 5ml. Utilizing crude plant extracts with well-established bioactive principles can help mitigate the harmful effects that synthetic nematodes have on bio-nature and plants. Tannin leaf and seed extracts, which were proven to be active in this investigation, may be useful in the development of novel nematicidal agents

Keywords: *Meloidogyne incognita*, Mortality, Okra yield, Physicinut, Root-gall nematode, Tannins

Introduction

The Global awareness on the hazards associated with chemical induced plant protection to man and the ecosystem cannot be over emphasized. A 2004 joint report from the World Health Organization (WHO), the United Nations Environment Programme (UNEP), and the Food and Agriculture Organization of the United Nations (FAO) revealed that a broad estimate of one million and five million cases of pesticide poisoning occur each year. More worrisome is the fact that, although, 25 % of the world's production of pesticides, is utilized by developing countries they experience 99 % of the death that occur due to pesticide poisoning (SP-IPM, 2000)

The most efficient way to manage high levels of *Meloidogyne* spp. in various farms is mainly through the use of chemical nematicides in control programs. However, several countries throughout the world have outlawed the use of such nematicides, which have methyl bromide and other dangerous substances as their active components (Onkendi *et al.*, 2014). The use of chemical methods to control these *Meloidogyne* spp. has drawbacks, including the fact that some of them are toxic to humans due to residues in the food chain, expensive for small-scale farmers, contribute to environmental pollution by destroying beneficial non-target organisms, and may eventually cause some resistance to the target nematode species.

In a comprehensive field survey on the occurrence, virulence, severity and root-knot nematode spread on vegetables, okra was highest in field infestation when compared to eggplant, tomato, cucumber, chilies, beans and cucurbits (Tariq-Khan, 2020). Historically, management of crop damage due to nematode attack has been achieved with the utilization of resistant variety, crop rotation, cultural measure practices and synthetic chemical nematicides which for a long time have been the most common approach (Siji *et al.*, 2010). This clearly brings to fore the urgent need for and importance of alternate management strategies which are ecologically friendly and potent against root-knot nematodes

Several plants have been evaluated for novel compounds with nematicidal potential against parasitic nematodes (Chitwood, 2002). Akinmoladun (2007) reported that the medicinal value of any plant is a function of their phytochemical constituents. The physicnut plant is one of those plants with abundance of phytochemicals (Devappa *et al.*, 2010). Igbinosa *et al.*, (2009) also reported that *J. curcas* contains biologically active phytochemicals amongst which are tannins.

Although, tannins have been shown to have anthelmintic attributes, particularly against nematodes of the gastrointestinal tract in ruminants, little is known on how well they would work as a substitute for conventional chemical nematicides in the management of okra root-knot nematode disease. Thus, the purpose of this investigation is to assess the effects of physicnut plant parts and tannins on okra root-gall nematodes *M. incognita* in both in vitro and in vivo settings.

Materials and Methods

Experimental Site

Experiments were conducted in the screen house of the Department of Crop Science and Technology, Federal University of Technology, Owerri, Nigeria and located on Lat 5° 27' 50.23" North and Longitude 07° 02' 49.33" East, at 55m above mean sea level. The soil was infested with *M. incognita* and classified as loamy sand (Eisenback *et al.*, 1981; Agu, 2008). The experiments were conducted twice, and the mean taken for most parameters since data collected were similar.

Plant Materials and Tannins Isolation

Fresh *Jatropha curcas* L. plant materials (leaves, roots, and seeds) were collected at the Teaching and Research Farm of Crop Science and Tech. Department (FUTO). After being sun-dried, they were ground into a coarse powder with a clean mortar and pestle, sieved to remove the fine particles, and kept in a dry location until needed. The purification process was mainly carried out in accordance with Strumeyer and Malin's (1975) procedure. Using an ultrasonicator, extracts of the dried ground plant parts (20 g) were prepared in 400 ml of 70 % aqueous acetone with 0.1 % ascorbic acid. To remove acetone, the supernatant that was produced after centrifuging for 15 minutes at 2500 rpm was evaporated under vacuum at a temperature of about 30 °C. The aqueous sample underwent lyophilization. Eighty percent aqueous methanol with 0.1 percent ascorbic acid was used to dissolve the dried material. After filtering the lyophilized material, a Sephadex-20 slurry was created by adding 300 ml of 80 % aqueous methanol to a chromatographic column containing 25 grams of Sephadex LH-20 and 0.1% of ascorbic acid. The filtrate was added to the column, and a conical flask was used to collect the eluted sample free of tannins. The sephadex LH-20 retained the tannins, which gave it a brownish colour. By adding 100 ml of 5 % aqueous acetone containing 1 mg/ml ascorbic acid to the column, the tannins were eluted.

The amount of ascorbic acid added during the isolation process was determined as Xmg of ascorbic acid, and the volume of the eluted tannins was measured in milliliters. By using a rotary vacuum evaporator, at 30 °C under vacuum, extraction of the acetone was carried out. The pure tannin-containing aqueous solution was then lyophilized in a beaker that had been previously weighed (W1) to determine the weight of the beaker plus the tannin solution (W2). The weight of the tannins containing ascorbic acid (Y, mg) was found by subtracting to get the weight difference of W2 and W1. Thus, the weight of tannins was calculated by subtracting (Y, mg - X, mg). However, 80 % aqueous methanol was used to obtain the final weight of tannins, which was devoid of ascorbic acid.

Test for Tannins

A yellow precipitate was formed on addition of three drops of 1 % lead acetate to 5 ml of the extract, indicating the presence of tannins (Ukoha *et al.*, 2011)

Nematicidal Assay

The effectiveness of crude tannins isolated from plant parts was assessed *in vitro* at 0, 5 and 10 ml against 30 nematode juveniles (J2s) after 24, 48, and 72 hours of incubation. The Petri dishes used were stored at room temperature ($\pm 30^{\circ}\text{C}$) on the laboratory bench. After being exposed to each treatment for 24, 48, and 72 hours, dead and live juveniles (J2) were counted using an electronic stereomicroscope at a magnification of 100X. When the juvenile (J2) did not respond to a fine needle used as a physical stimulus, it was deemed dead (Hong *et al.*, 2007). The ratio of the number of juvenile deaths to the total number of juveniles was used to compute percentage mortality. For instance, % Mortality = $X/N \times 100/1$, where X is the number of juvenile deaths (J2). N = Initial Juvenile Number (J2).

Screen House Experiment

The study was carried out using a 3x6 factorial experiment placed in a Completely Randomized Design. The physiconut had three levels of treatment (root, seed, and leaves), while the tannins had six; [0.00 (control), 1, 2, 3, 4, and 5.00 ml]. Consequently, 18 treatment combinations were allocated randomly across 90 pots in the screen house, with each containing 4 kg of steam-sterilized soil and replicated 5 times

Nematode Inoculum

The inoculum came from Owerri population of *M. incognita* that was maintained for up to eight weeks on susceptible okra (NH47-4). The infected (galled) portion was recovered by upturning the pots to release the root system. Additionally, chopped galled roots were added to a warren blender along with one liter of water. The blending was only allowed to run for 3-second intervals to prevent inactivating the nematode larvae. More water was added to the slurry to bring it up to 1200 ml. A Petri-dish was filled with thirty milliliters (30 ml) of this mixture, and the larvae were counted under a stereomicroscope. Three counts yielded an average of 430 ± 1.50 larvae.

Seed Sowing

The National Horticultural Research Institute, Okigwe in Imo State, Nigeria, provided the test okra cultivar that was utilized. However, surface sterilization was carried out on test okra cultivar (NH47-4) before planting. Commercial sodium hypochlorite solution (NaOCL) was used to soak okra seeds and later washed thoroughly in clean water. In each potted soil, two sprouting seeds were sown and before inoculation, the seeds were thinned to just one per pot.

Inoculation and Phytochemical application

Okra cultivar was inoculated with 100mL (about 1200 infective nematode larvae) of the inoculum at the base of each potted plant 10 days after planting. Using the formula $V_1C_1 = V_2C_2$, (Naz *et al.*, 2013) the tannins extract was diluted in distilled water to reach a final concentration of 50 mg/ml. Seven days following inoculation, this was administered at the base of the potted plants five days later at 0, 1, 2, 3, 4, and 5.00 ml per plant. Plants that were inoculated but not treated with tannins acted as the control group. Plants inoculated were housed in a screen house with good ventilation. The soil's average temperature was between 30 and 32°C

Growth and Yield Measurement

A Brookfield RVT viscometer (manufactured in the United States by Brookfield Engineering Laboratories) was used to measure and read the mucilaginous characteristic of okra fruits with No. 1 spindle at 15 r.p.m. (15 revolutions per minute). This procedure was repeated three times and the mean taken for each treatment (Sopade *et al.*, 1992). Other data collected included leaf areas, number of leaves, plant heights, 100 dry seed weights and fresh pod weights.

Infection Assessment

Ten weeks after inoculation when data on growth and yield attributes had been collected, the roots were checked for infections. By flipping the pots, root systems were freed and retrieved undamaged before washing with water to get rid of soils stuck to the roots. Agu and Ogbuji (1996) were used to score root systems, with 0 denoting absence of infection (no galls present), 1 = Infrequent infection or infection that is rare (1-3 galls visible), 2 = Mild infection (4-10 galls in place), 3 = Moderate infection with 11-30 galls visible and 4 = Serious infection (with 30 or more galls).

Statistical Analysis

Using the software GENSTAT (Edition 4), data collected was subjected to analysis of variance. Fisher's Least-Significant difference (F - LSD) was used to separate differences between means at the 5 % probability level. Root-gall index and yields were correlated using correlation analysis with SPSS version 22.0.

Results

Results of the study showed that the percentage yields of tannins isolated from *J. curcas* plant were as follows; seed (49.05 %), leaf (11.04 %) and root (12.65 %). The reaction of nematode juveniles (J2) to tannins extract of *J. curcas* seeds, leaves and roots are shown in Figure 1. Death of most nematodes (J2) i.e. (28/30) was observed to have significantly ($P < 0.05$) occurred on application of *J. curcas* Tannins. Nematode mortality was highest (94%) at 10 ml of tannin treatment compared with the controls (Juveniles in distilled water), and this differed statistically ($P < 0.05$) with mortality obtained with 5 ml of tannins application (74%). Juvenile mortality was found to have risen at higher rates of tannins applied

Most nematode (J2) mortality was observed when the juveniles were exposed to tannins at 72-hour intervals, as shown in the interactions with the tannins treatment's hour of exposure. Nematode mortality at 48-hour exposure intervals was next. However, Intervals of 24 hours saw the least number of nematode mortalities. Consequently, it was found that the number of hours that the nematode (J2) was exposed to the tannin treatment increased along with its mortality. Significant increases in nematode mortality were seen on application of 5 and 10 ml of plant extract containing *Jatropha* tannins. Applying root tannins caused the greatest nematode mortality (98%). This was followed by seed tannins (90 %) and leaf tannins (81 %) respectively.

Figure 2 shows how tannins extracted from different plant parts affected the degree of root-gall nematode infection on okra. Root-gall infection assessment on all okra plants in the inoculated untreated (control) revealed a gall index of 4 (severely galled). However, a progressive reduction in gall index was noticed with corresponding increase in tannins treatments applied. At 5 ml application of leaf and seed tannins lowest gall index (1.00) was observed in the test okra plants which indicates that the roots were rarely galled.

Table 1 displays the impact of tannins and *Jatropha* parts on height of okra, leaf area, pod weight, and 100 dry seed weight as affected by root-knot nematode. The plants on pots treated with various rates of tannins extracts had reduced root galls and produced significantly ($p < 0.05$) higher plant heights than the severely galled untreated plants. This was particularly so with the interactions of *Jatropha* parts and tannins where 5 ml of leaf tannins produced the highest plant heights. Subsequently plants treated with 5 ml of seed tannins came next which significantly ($p < 0.05$) differed from root tannins. Again, both tannins and *Jatropha* parts had significant ($p < 0.05$) effect on okra leaf area. Although interactions were not significant, highest leaf area was produced on plants treated with 5 ml of leaf tannins which significantly ($p < 0.05$) differed from seed and root tannins.

With increased application of tannin treatments, mean pod weights of okra were shown to have increased with fewer root-galls. When okra plants were treated with 5 mL of leaf and seed tannins, their pod weights were the highest and differed from the untreated control group and those treated with root tannins by a significant ($p < 0.05$) margin. When compared to other rates, the yield of 100 dry seed weights was highest when *Jatropha* root tannins were applied at 5 ml. Plants that received seed tannins treatment came next. The lowest 100 dry seed weights were found in plants that received leaf tannins treatment.

The impact of Jatropha ¹ tannins and parts on okra's mucilaginous quality as affected by root-gall nematode is illustrated in Figure 3. When tannins were administered at higher rates, the mucilaginous property substantially improved. In comparison with leaf and seed tannins, root tannins treatment exhibited significantly ($p < 0.05$) greater mucilaginous properties. When comparing plants with the highest mucilaginous property to other rates and the untreated control, root tannins at 5 ml significantly ($p < 0.05$) increased the mucilaginous property (2124 mpa*s) of the plants. Mucilaginous property of okra plants treated with 5 ml of leaf (1605 mpa*s) and seed (2105 mpa*s) tannins followed next before the untreated control (1000 mpa*s)

¹³ Correlation analysis ³⁷ the root-gall index and yield attributes of okra as influenced by the tannin's extracts are presented in Table 2. The results showed that the root gall index had a negative and highly significant correlation with the growth and yield attributes. However, mucilaginous property correlated positively and significantly with plant height. The same was true for 100 dry seed weights, pod weights and leaf areas





Figure 1: Relationship between *J. curcas* Tannins, Plant Parts, Hour of Exposure and Mortality Effect on *M. incognita*



Scale of rating: 0 = not infected (no galls), 1 = Seldomly infected (1-3 galls), 2 = Mild infection (4-10 galls), 3 indicates a moderate infection (between 11 and 30 galls), and 4 indicates a severe condition or infection (with over 30 galls present).

Figure 2: Effect of *Jatropha* Tannins and plant parts on root-gall nematode (*M. incognita*) infection

Table 1: Effect of *Jatropha curcas* Tannins and plant parts on Okra plant heights, Leaf areas, Pod weights and 100 dry seed weights as affected by root-gall nematode

Tannins (ml)	Plant Heights (cm)			Leaf Areas (cm ²)			Pod weights (g)			100 dry seed weights (g)						
	Jatropha Parts			Jatropha Parts			Jatropha Parts			Jatropha Parts						
	Leaf	Root	Seed	Mean	Leaf	Root	Seed	Mean	Leaf	Root	Seed	Mean	Leaf	Root	Seed	Mean
Untreated (Control)	9.92	9.59	9.51	9.67	10.46	9.79	11.17	10.47	1.90	1.70	1.30	1.63	2.70	2.90	1.90	2.50
1	10.12	10.04	11.28	10.48	11.09	11.46	12.93	11.83	2.30	2.00	2.40	2.23	3.00	3.33	3.50	3.26
2	10.77	11.39	12.67	11.61	12.38	12.74	14.06	13.06	2.40	1.85	3.60	2.61	2.85	3.40	4.50	3.58
3	14.81	11.99	13.77	13.52	21.45	15.29	15.54	17.43	4.06	2.80	3.90	3.58	3.80	5.06	4.80	4.55
4	15.87	13.21	15.36	14.81	25.15	18.03	18.88	20.69	4.43	2.84	4.29	3.85	3.83	5.42	5.28	4.84
5	20.79	14.62	17.59	17.67	34.04	24.12	27.48	28.55	4.95	3.46	4.80	4.40	4.30	6.91	5.80	5.67
Mean	13.71	11.8	13.36		19.1	15.24	16.68		3.34	2.44	3.38		3.41	4.49	4.29	
LSD _{0.05} (Jatropha parts)	0.93				2.78				0.07				0.07			
LSD _{0.05} (Tannins)	1.31				3.93				0.10				0.10			
LSD _{0.05} (Jatropha parts X Tannins)	ns				ns				0.17				0.17			

ns = Not Significant,




Figure 3: Effect of *Jatropha* Tannins and plant parts on Okra's mucilaginous property as affected by root-gall nematode

Table 2. A correlation matrix illustrating the effects tannins treatment on plant characteristics and the root-gall index

1 st Trial	PH	LA	NL	PW	HSW	MP	RGI
PLHT	1	.771**	.616**	.814**	.533**	.608**	-.738**
LFA		1	.553**	.710**	.490**	.597**	-.676**
NOLVS			1	.664**	.479**	.510**	-.626**
PODWT				1	.653**	.615**	-.847**
HUNSEDWT					1	.870**	-.583**
MUCIL						1	-.623**
RGI							1
2nd Trial							
PH	1						
LA	.785**	1					
NL	.639**	.584**	1				
PW	.820**	.728**	.616**	1			
HSW	.550**	.515**	.465**	.662**	1		
MP	.586**	.588**	.397**	.606**	.860**	1	
RGI	-.682**	-.638**	-.488**	-.808**	-.588**	-.616**	1

LA = Leaf Areas

PH = Plant Heights

NL = Number of Leaves

HSW = 100 Seed Weights

MP = Mucilaginous Property -mpa*s

RGI = Root-Gall Index

Discussion

The percentage yield of tannins shows that its availability differs among the *J. curcas* parts with the highest concentration in the seed, followed by the root and leaf respectively. The nematotoxic impact of *J. curcas* tannins may have contributed to the nematode juvenile mortality observed in comparison to the control. Molan (2014) confirmed that condensed tannins inhibited egg hatching, slowed down larval development and killed undeveloped larvae thereby disrupting the life cycle of the nematode *Teladorsagia circumcincta*. Corroboratively, Tong (2022) stated that the direct mechanisms involve restricting larval development to reduce the establishment of infected third stage larvae (J3) in the host and decreasing spawning to inhibit the motion performance of the parasite.

Humaira *et al.*, (2020) found that the concentration of plant extract has a significant impact on larval mortality. The higher nematode mortality observed as a result of the equivalent increase in tannin application rate is in line with these findings. With more hours of tannins treatment exposure, there was a noticeable rise in nematode juvenile mortality. In a similar vein, extracts of *Acacia modesta* plant parts showed extremely encouraging mortality of greater than 60 % after 72 hours of treatment, according to Humaira *et al.* (2020).

All untreated inoculated okra plants were observed to be severely galled due to root-gall nematode activity on the root. Castillo *et al.*, 2001 reported that the conspicuous giant cells (galls) produced on infected roots and excessive lateral root growth is the major symptoms of root-knot nematode infection which decrease the ability of infected plants to translocate nutrient and water from soil to vegetative organs of plants. Tannins extract of various plant parts significantly reduced root-gall indexes. This performance can be seen more with the application of leaf and seed extracts of tannings. The performance of the seed m

Tannin extracts applied to okra plants showed improved yield performances and fewer galls than the heavily galled untreated plants which served as the control. This might have been due to the nematocidal action of tannins. Among the naturally occurring substances derived from plants, tannins have been found shown to have anthelmintic qualities, particularly for ruminant gastrointestinal nematodes. Additionally, they are antagonistic to several bacteria, yeasts, and fungi. Maistrello *et al.*, (2010) further reported that tannins are substances that may contribute to the plant's passive defense by acting as chemical barriers to the parasite's entry into the roots and may strengthen the resistance of the host to nematode infection.

Tannins nematocidal action, inhibited galling and promoted growth. This may be the reason for the higher plant heights in the treated plants compared to the untreated control after 5 ml of leaf and root tannins were applied. Agaf findings of Maistrello *et al.*, (2010), also confirming that greatest doses of tannins administered at transplant and again two weeks later considerably decreased the root-gall index when compared to the inoculated and untreated control, are consistent with this. The reduction in plant heights in the severely galled untreated control plants could be as a result of the incidence and severity of the nematodes on the inoculated control plants. Mukhtar *et al.* 2013 reported that all inoculums' levels of *M. incognita* caused significant reductions in plant heights at all plant ages of okra.

There were also observed increases recorded in the leaf area on application of 5 ml of Leaf Tannins in the treated than the untreated plants. This enhanced growth could be as a result of the suppressive and lethal effects of the leaf Tannins extracts. Azhagumurugan. and Rajan, (2014) reported that extracts from plant leaf are characterized by their lipophilic properties that enable them dissolve the cytoplasmic membrane nematodes. This might have led to improved physiological activities of the plant. Again, the production of leaves increased with increased rates/dosage of tannins treatments.

Reduction in leaf areas of plants in the infected but untreated control pots may have been caused by reduction in the chlorophyll content of the infected plants. According to Parveen *et al.* (2006), a decrease in the chlorophyll content of the infected plants may have contributed to the reduction in leaf areas of the plants in the infected but untreated control pots. Furthermore, infection with *M. incognita* also resulted in a decrease in leaf area, as reported by Jonathan and Rajendan (2002).

Comparing okra plants treated with 5 mL of leaf and seed tannins to plants treated with root tannins, the former yielded the largest pod weights. According to Ojiako *et al.* (2015), the repellency activity of the seed extract was in fact superior to that of root extracts, regardless of the solvent utilized. Khan *et al.*, (2017) have also reported on the effectiveness of plant leaf extract. The treated okra plants yielded fruits with a higher fruit weight than the untreated ones. The fruits with the lowest weight were the untreated (controls). This indicates that the fruits on the treated plants were significantly heavier than those of the untreated plants. This is consistent with research by Mbah *et al.* (2005), who discovered that tomato plants treated with plant leaf extracts produced a higher yield than plants left untreated (control).

Tannin treatments suppressed nematode activity, which made it possible for the plants to take up nutrients for healthy growth and development. The increased okra pod weights recorded could therefore, be attributed to improved physiological activities of the Tannins treated plants. According to Agu *et al.* (2013), higher leaf production on tomato plants under root-gall nematode control boosted the plants' rate of photosynthetic activity and thus produced higher fruit harvests. Fruit weights are significant since they are a key factor in determining market values. The recorded improvement on 100 dry seed weights was as a result of decreases in root-gall damage by the nematicidal effect of tannins, thereby improving nutrient absorption and translocation to vegetative organs. Agu *et al.* (2013) stated that plants with fewer root-galls would translocate more nutrients to vegetative organs for proper seed and pod development than heavily galled roots.

An increase in the rate at which tannins was applied was also found to enhance the mucilaginous quality of okra pods. Agu *et al.*, (2009) also observed a continuous increase in the mucilaginous characteristic of pods as galling response to *M. incognita* diminished with greater carbofuran treatment. Furthermore, he also reported a continuous reduction in galling as *J. curcas* application increased.

Growth and yield quality of okra reduced with an increase in gall count, in accordance with the observed negative association between the root-gall index and okra yields. Anon and Mckenry (2010) suggested that this might be caused by aberrant cells called galls that interfere with the movement of nutrients and moisture within the root system of plants. Higher photosynthesis and improved growth and yield qualities of okra were attributed to higher numbers of leaves and areas

of leaves at decreased galling responses, as demonstrated by the positive correlation among leaf area, number of leaves produced, pod weights, 100 dry seed weights, and mucilaginous property.

Conclusion

Tannin's extract of *J. curcas* has been shown to be lethal to the root-knot nematode *M. incognita* in this study. The error bars have underscored the statistical significance of these results, highlighting the importance of Tannins rate and exposure duration, along with the specific *Jatropha* part used in determining the mortality rate of *M. incognita*. The development of new nematicidal agents may benefit from the use of leaf, seeds and even root extracts of tannins, which were found to be active in this study. As a result, there is hope for the identification of new classes of nematicides from natural plants to replace the synthetic nematicides currently in use. Present research findings suggest that integrated disease management programs can effectively deploy crude seed and leaf tannins extracts against root-knot nematodes at appropriate concentration and rate reported in this study, due to their high nematicidal potential. Furthermore, the study's findings on the efficacy of *Jatropha* tannins present enormous opportunities for novel biotechnological applications in nematode formulations. However, to build on the findings of this study, examination in a field setting is recommended.

Acknowledgement

The first author acknowledges the study grant from Federal University of Technology, Owerri, Nigeria, which enabled the sponsorship of this study. The authors also wish to acknowledge the support of Prof. C.M. Agu for his immense contribution towards the conception and technical assistance offered in this work.

Author contribution OEV, and OOF planned the experiments, OEV, EN, NE, and EOC statistically interpreted the result, MAO, ES, KB AND UQC made the write up and OEV, MAO and OOF statistically analyzed the data and made illustrations

Conflict of Interest

All authors have no conflict of interest about the results presented in this article

References

- Agu, C.M and R.O. Ogbuji. 1996. Soybean resistance to the root-gall nematode, as influenced by Potassium nutrition. *East African Agricultural Journal* 66(3), 273-276.
<https://www.africabib.org/rec.php?RID=Q00049962>
- Agu, C.M., C.O. Mbuka and V.E. Ogwudire. 2009. Influence of root-gall nematode on mucilaginous property and nutrient composition of Okra (*Abelmoschus esculentus* L. Moench). *Bionature*, **29**(2), 73-77.
<https://globalpresshub.com/index.php/BN/article/view/447>
- Agu, C.M., A.C. Nwagbaga, C.G. Uwandu, P. Onoh, C.A. Mbuka, F.O. Ojiako, N.K. Ndulue and V.I Ogwudire. 2013. Effect of *Jatropha Curcas* on Root-gall Nematode Disease and the Consequent Proximate Composition of Okra. *Bionature Journal of India*, **33** (2), 59-62. <https://globalpresshub.com/index.php/BN/article/view/562>
- Akinmoladun, A.C., E.O. Ibukun, E. Afor, E.M. Obuotor and E.O Farombi. 2007. Phytochemical Constituents and Antioxidant Activity of Extract from the Leaves of the *Ocimum gratissimum*. *Sci. Res. Essay*, **2**, 163-166.
<http://www.academicjournals.org/SRE>.
- Anwar, S.A., M.V and McKenry. 2010. Incidence and reproduction of *Meloidogyne incognita* on vegetable crop genotypes. *Pakistan Journal of zoology*, 42(2):135-141.
[http://zsp.com.pk/pdf/135-141%20\(5\).pdf](http://zsp.com.pk/pdf/135-141%20(5).pdf)
- Azhagumurugan., C and M.K. Rajan. 2014. Extract of Nilakumil, (*gmelina asiatica*) against the Root knot Nematode, (*Meloidogyne incognita*). *Research Journal of Recent Sciences*. Vol. 3(ISC-2013), 264-266 ISSN 2277-2502. www.isca.in.
- Castillo, P.D.I., M. Vito, N. Vovlas and R.M. Jiménez-Díaz. 2001. Host-parasite relationships in root-knot disease of white mulberry. *Plant Disease*, 85, 277-281.
- Chitwood, D.J. 2002. Phytochemical Based Strategies for Nematode Control. *Annu. Rev. Phytopathol.* 40:221–49. doi: 10.1146/annurev.phyto.40.032602.130045
- Devappa, R.K., H.P.S. Makkar and K. Becker. 2010. *Jatropha* toxicity- a review *Journal of Toxicology and Environmental Health, Part B*, 13, pp. 476-507. doi: 10.1080/10937404.2010.499736.
- Eisenback, J.D., H. Hirschmann, J.N. Sasser and A.C. Triantaphyllou. 1981. A guide to the four most common species of root-knot nematodes (*Meloidogyne* spp) with a pictorial key. *A cooperative publication of North Carolina State University and USAID.*, pp: 48.
https://pdf.usaid.gov/pdf_docs/pnaaq221.pdf
- Hong, L., G. Li, W. Zhou, X. Wang and K. Zhang. 2007. Screening and isolation of a nematicidal sesquiterpene from *Magnolia grandiflora*. *Pest Managment. Science*. 63, 301-305. doi: 10.1002/ps.1337.

- Humaira – Shad, A.A., D. Muhammad and H.U. Shah. 2020. Evaluation of Some Medicinal Plant Extracts for Their Nematicidal Properties Against Root-Knot Nematode, *Meloidogyne Incognita*. *Applied Ecology and Environmental Research* 18(2):2475-2482. <http://www.aloki.hu>. ISSN 1589 1623 (Print). ISSN1785 0037 (Online) DOI: http://dx.doi.org/10.15666/aeer/1802_24752482
- Igbinosa, O.O., E.O. Igbinosa and O.A. Aiyegoro. 2009. Antimicrobial Activity and Phytochemical Screening of Stem Bark Extracts from *Jatropha curcas* (Linn). *African Journal of Pharmacy and Pharmacology*, **3** (9), 58-62. www.internationalscholarsjournals.org
- Janathan. E.I and G. Rajendran. 2002. Assessment of avoidable yield loss in banana due to root-knot nematode *Meloidogyne incognita*. *Indian J. Nematol.*, **30**: 162-164. <https://www.cabdirect.org/cabdirect/abstract/20023059279>
- Khan, A., M. Asif, M. Tariq, B. Rehman, K. Parihar and M.A. Siddiqui. 2017. Phytochemical investigation, nematostatic and nematicidal potential of weeds extract against the root-knot nematode, *Meloidogyne incognita* in vitro. *Asian Journal of Biological. Science*. **10**: 38-46.
- Maistrello, L., Vaccari, G and N. Sasanelli. 2010. Effect of chestnut tannins on the root-knot nematode *Meloidogyne javanica* *Parasitological Institute of SAS, Helminthologia*, **47**, 1: 48 – 57. DOI 10.2478/s11687-010-0008-9
- Mba, O.I., N.C. Ohazuruike and E.A. Okorochoa. 2005. An assessment of the nematicidal potential of some plant products in the control of root-knot nematode (*M. incognita*, Kofoid and White) on tomato plant. *Proceedings of the 32nd Annual Conference of Nigeria Society for Plant Protection. 15th – 20th May 2005. University of Port Harcourt, Nigeria*, 37-41.
- Molan, A.L. 2014. Effect of purified condensed tannins from pine bark on larval motility, egg hatching and larval development of *Teladorsagia circumcincta* and *Trichostrongylus colubriformis* (Nematoda: Trichostrongylidae). *Folia Parasitol (Praha)*. **61**:371–6. doi: 10.14411/fp.2014.036.
- Mukhtar, T., A. Israr, M. Zameer, H.M. Arshad, S.B. Kayani, R.A. Murad and A. Muhammad. 2013. Estimation of Damage to Okra by Root-Knot Disease Mated by *Meloidogyne incognita*. *Pakistan Journal of Botany*, **45** (3): 1023 – 1027. <https://www.researchgate.net/publication/236975524>
- Naz, I., Saifullah, M.R and Khan. 2013. Nematicidal activity of nonacosane-10-ol and 23a-Homostigmast-5-en-3 β -ol isolated from the roots of *Fumaria parviflora* (Fumariaceae). *Journal of Agricultural and Food Chemistry*, vol. 61, no. 24, pp. 5689-5695. <http://dx.doi.org/10.1021/jf401309r>. PMID:23713689.
- Ojiako, F.O., G.O. Ihejirika and U.O. Aguwa. 2015. Comparative bioactivity of different solvent extracts of the root and seeds of *Jatropha curcas* L. and Chlorpyrifos, against tailor ants (*Oecophylla longinoda* Latreille) (Hymenoptera: Formicidae). *FUTO Journal Series*

(*FUTOJNLS*).1 (2); p-ISSN: 2467-8325; e-ISSN: 2476-8456.
https://www.researchgate.net/publication/298972376_

- Onkendi, E.M., G.M. Kariuki, M. Mara and L.N. Moleleki. 2014. The threat of root-knot nematodes (*Meloidogyne* spp.) in Africa: a review. *Forestry Agriculture and Biotechnology Institute (FABI)*. Department of Microbiology and Plant Pathology. University of Pretoria. South Africa. doi:10.1111/ppa.12202
- Parveen, K., A. Haseeb and P.K.I. Shukla. 2006. Pathogenic potential of *Meloidogyne incognita* on *Mentha arvensis* cv. Gomti. *Ind. J. Nematol.*, 36: 177-180.
<https://www.indianjournals.com/ijor.aspx?target=ijor:ijn&volume=36&issue=2&article=001>
- Siji, J.V., C.A. Jayaprakas, M.S. Sheela and C. Mohandas. 2010. Efficacy of *Cleome viscosa* L. against *Meloidogyne incognita* infestation in okra (*Abelmoschus esculentus* L.). *Thai Journal of Agricultural Science* 43 (3): 151-156. <https://www.thaiscience.info/Journals/Article/TJAS/10656559.pdf>
- SP-IPM, 2000. Biological alternatives to harmful chemical pesticide. *IPM Research brief no. 4*. The Systemwide Program on Integrated Pest Management for improved livelihood. www.spipm.cgiar.org.
- Sopade, P., Ajisegiri, E and Badau. 1992. The Use of Peleg's Equation to Model Water Absorption in some Cereal Grains during Soaking. *Journal of Food Engineering* 15:269-283.
- Strumeyer, D.H and M.J. Malin. 1975. Condensed Tannins in grain sorghum Isolation, Fraction, and Characterization. *Journal of Agricultural and Food Chemistry*. 23 (5): 909-914. doi:10.1021/jf60201a019
- Tariq-Khan, M. S.Z.A. Gardazi, A.D.A. Khan, M. Iyas and M. Ahmad. 2020. Virulence and distribution trends of root-knot nematode (RKN) fauna on summer vegetables in District Bagh, Azad Jammu and Kashmir (Pakistan). *Pakistan Journal of Nematology*, 38(2): 139-148. doi <http://dx.doi.org/10.17582/journal.pjn/2020.38.2.139.148>
- Tong Z, W He, X Fan, A Guo, (2022) Biological Function of Plant Tannins and Its Application in Animal Health. *Front. Vet. Sci.* 8:803657. doi: 10.3389/fvets.2021.803657.
- Ukoha PO, AC Egbuonu, Cemaluk, L Obasi, Nnamdi, PM Ejikeme (2011). Tannins and other phytochemical of the *Samanea saman* pods and their antimicrobial activities. *African Journal of Pure and Applied Chemistry* Vol. 5(8), pp. 237-244, <http://www.academicjournals.org/AJPAC> ISSN 1996 - 0840.

The Role of Tannins as Plant Based Nematicide in Root-Gall Nematode Management of Okra (*Abelmoschus esculentus*) for Improved Yield

ORIGINALITY REPORT

17%

SIMILARITY INDEX

13%

INTERNET SOURCES

9%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1	bioscientistjournal.com Internet Source	4%
2	Submitted to Federal University of Technology Student Paper	1%
3	www.aujst.com Internet Source	1%
4	www.researchgate.net Internet Source	1%
5	oer.unn.edu.ng Internet Source	1%
6	www.ncbi.nlm.nih.gov Internet Source	1%
7	Nwakuche Chinenye Onwubiko. " Genetic diversity in Bambara groundnut { (L.) Verdc.} ", Agricultura Tropica et Subtropica, 2021 Publication	<1%
8	www.giikorea.co.kr Internet Source	

<1 %

9

CM Agu. "Effect of phosphorus fertilizer on root-knot nematode disease of soyabean", Tropical Science, 2006

Publication

<1 %

10

Wang Kui, Luo Chao, Liu Hao, Xu Jianmei, Sun Weibo, Zhou Ligang. "Nematicidal activity of the alkaloids from *Macleaya cordata* against certain nematodes", African Journal of Agricultural Research, 2012

Publication

<1 %

11

Aladesanwa, R.D.. "Screenhouse evaluation of atrazine for soil residual activity on growth, development and nutritional quality of okra (*Abelmoschus esculentus* Moench) in southwestern Nigeria", Crop Protection, 200510

Publication

<1 %

12

www.isca.isca.in

Internet Source

<1 %

13

J. I. Oluwatayo, C. I. Jidere, A. Nwankiti. "Nematicidal Effect of Some Botanical Extracts for the Management of *Meloidogyne incognita* and on Growth of Tomato", Asian Journal of Agricultural and Horticultural Research, 2019

Publication

<1 %

14

L. Maistrello, G. Vaccari, N. Sasanelli. "Effect of chestnut tannins on the root-knot nematode *Meloidogyne javanica*", *Helminthologia*, 2010

Publication

<1 %

15

www.scielo.br

Internet Source

<1 %

16

C. M. Agu. "Effect of Urea Fertilizer on Root-Gall Disease of *Meloidogyne javanica* in Soybean (*Glycine max* (L.) Merrill)", *Journal of Sustainable Agriculture*, 2002

Publication

<1 %

17

E.E. Itoandon, S.O.A. Olatope, O.O. Shobowale. "Preliminary Phytochemical Analysis and Antimicrobial Properties of Crude Extract of *Combretodendron macrocarpum* Stem Bark", *Nigerian Food Journal*, 2012

Publication

<1 %

18

Grace Mohoanatsso Tefu, Dirk De Waele, Hendrika Fourie, Mieke Stefanie Daneel. "Nematode-suppressive and plant growth-enhancing effects of spp. derivatives against on tomato ", *International Journal of Pest Management*, 2024

Publication

<1 %

19

academicjournals.org

Internet Source

<1 %

20

"PHYTOCHEMICAL BASED STRATEGIES FOR NEMATODE CONTROL1", Annual Review of Phytopathology, 09/2002

Publication

<1 %

21

Faryad Khan, Mohd Asif, Amir Khan, Moh Tariq, Taruba Ansari, Mohammad Shariq, Mansoor Ahmad Siddiqui. "Evaluation of the nematicidal potential of some botanicals against root-knot nematode, *Meloidogyne incognita* infected carrot: In vitro and greenhouse study", Current Plant Biology, 2019

Publication

<1 %

22

d197for5662m48.cloudfront.net

Internet Source

<1 %

23

www.iosrjournals.org

Internet Source

<1 %

24

A.M. Babu, Jalaja S. Kumar, P.K. Mishra, B. Dayakar Yadav, S.M.H. Qadri. " Root-knot development in mulberry infested with ", Archives Of Phytopathology And Plant Protection, 2012

Publication

<1 %

25

S. Cherian. "Cadmium uptake and distribution in tomato plants (*Lycopersicon esculentum*

<1 %

Mill)", The South Pacific Journal of Natural Science, 2007

Publication

26

epdf.pub

Internet Source

<1 %

27

pr.hec.gov.pk

Internet Source

<1 %

28

repository.nwu.ac.za

Internet Source

<1 %

29

www.syncsci.com

Internet Source

<1 %

30

Hadis Mostafanezhad, Navazollah Sahebani, Shaheen Nourinejhad Zarghani. " Control of root-knot nematode () with combination of and salicylic acid and study of some plant defense responses ", Biocontrol Science and Technology, 2013

Publication

<1 %

31

Matos Martins Diniz Guilherme, dos Santos Candido Willame, Henrique Costa Silva Edgard, Vinícius Marin Marcus et al. "Screening melon genotypes for resistance to Meloidogyne enterolobii", African Journal of Agricultural Research, 2016

Publication

<1 %

32

Parvatha P. Reddy. "Nematode Diseases of Crops and their Management", Springer

<1 %

Science and Business Media LLC, 2021

Publication

33

Plant Growth Promoting Rhizobacteria for Horticultural Crop Protection, 2014.

Publication

<1 %

34

Wantana Mongkolvisut, Somyote Sutthivaiyakit, Heiko Leutbecher, Sabine Mika et al. " Integerrimides A and B, Cyclic Heptapeptides from the Latex of ", Journal of Natural Products, 2006

Publication

<1 %

35

escijournals.net

Internet Source

<1 %

36

kobra.uni-kassel.de

Internet Source

<1 %

37

uniuyo.edu.ng

Internet Source

<1 %

38

www.fspublishers.org

Internet Source

<1 %

39

www.globalpresshub.com

Internet Source

<1 %

40

www.scialert.net

Internet Source

<1 %

41

C. M. Agu. " Control of Root-Knot Nematode Disease on Tomato by (Stapf & Hubb), (Lin)

<1 %

and (Pegs) ", Journal of Sustainable
Agriculture, 2005

Publication

42

R. Khan, I. Naz, S. Hussain, R. A. A. Khan, S. Ullah, M. U. Rashid, I. Siddique.

"Phytochemical management of root knot nematode (*Meloidogyne incognita*) kofoid and white chitwood by *Artemisia* spp. in tomato (*Lycopersicon esculentum* L.)", *Brazilian Journal of Biology*, 2019

Publication

<1 %

43

Waceke J. Wanjohi, Geoffrey O. Wafula, Charles M. Macharia. "Integrated Management of Fusarium Wilt-Root Knot Nematode Complex on Tomato in Central Highlands of Kenya", *Sustainable Agriculture Research*, 2018

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On