



Official publication of Pakistan Phytopathological Society
Pakistan Journal of Phytopathology

ISSN: 1019-763X (Print), 2305-0284 (Online)
<http://www.pakps.com>



GENETIC DIVERSITY AND NUCLEOTIDE SEQUENCE CHARACTERIZATION OF THE P4 GENE OF *RICE TUNGRO BACILLIFORM VIRUS (RTBV)* IN SOUTH KALIMANTAN, INDONESIA

Noor Aidawati*, Saipul Abbas, Elly Liestiany

Department of Plant Protection, Faculty of Agriculture, Lambung Mangkurat University. Jl.Jend. A. Yani Km.36 Banjarbaru. South Kalimantan 70714 Indonesia.

ABSTRACT

Rice tungro bacilliform virus (RTBV) is a serious threat to rice production in various regions in Indonesia including South Kalimantan. Cautious and flexible management strategies are of utmost importance. This study investigates the genetic diversity within the P4 gene of *RTBV* in South Kalimantan, Indonesia, to inform targeted disease management strategies. The PCR method was used to amplify P4 gene fragments from *RTBV* samples collected from 2 locations, namely *Karang Buah* Village and *Anjir Pasar Lama*. The PCR results successfully detected fragments of 430 bp, which were then analyzed in sequence to identify genetic variations. The characterization results showed that the presence of significant genetic diversity among *RTBV* isolates. These variants include differences in nucleotides at specific positions in the P4 gene, indicating the adaptation of the virus to local conditions. These findings provide a solid basis for designing more precise control strategies, such as the development of rice varieties that are more resistant to certain *RTBV* variants or the implementation of more specific management measures.

Keywords: PCR; P4 Gene; *RTBV*; Rice; Sequence Characterization.

INTRODUCTION

Rice tungro bacilliform virus (RTBV), a significant pathogen within the family *Caulimoviridae* and the genus *Badnavirus* (Bömer *et al.*, 2018), poses a critical threat to rice production in Indonesia, affecting both yield and quality. The virus has a single-stranded DNA-shaped genome and is generally transmitted via insect vectors. The transmission of *RTBV* by the leafhopper *Nephotettix vitrescens* is dependent upon the *Rice tungro spherical virus (RTSV)*. The transmission of *RTBV* and *RTSV* is semi-persistent in adult leafhoppers (Hibino and Cabauatan, 1987). In addition to an eye examination, scientific confirmation of the condition is currently achieved through the use of polymerase chain reaction (PCR) technologies that detect viral nucleic acids (Yee *et al.*, 2017). The PCR method is simpler and quicker

than other methods, it is highly beneficial for identifying the presence of rice viruses (Abbas *et al.*, 2020). PCR analysis was used to identify and amplify the P4 gene of *RTBV*, allowing detection of the presence of the virus in the sample. P3 and P4 gene amplification was done using PCR. The primer sequences used were CP5-5' - ATGAATACGGCCGCGGCTAGAAATC-3' and 5' - CTATTTTCGGGTTCTGGACCTGGCA-3' (Khanal and Ali, 2019; Pouraziz *et al.*, 2023)

Characterization of the nucleotide sequence of the P4 *RTBV* gene involves determining the nucleic acid sequence, thus providing an understanding of the structure and genetic variation of the virus. Reduced 21-nt siRNA levels and increased 22-nt siRNA levels were linked with *RTBV* protein P4's suppression of cell-to-cell spread of silencing and enhancement of cell-autonomous silencing (Rajeswaran *et al.*, 2014). The strategy of tungro disease control in rice plants is strengthened through the genetic understanding of *RTBV*, resulting from PCR analysis and P4 gene sequencing. A key component of disease control

Submitted: December 20, 2023

Revised: February 17, 2024

Accepted for Publication: May 25, 2024

* Corresponding Author:

Email: nooraidawati@ulm.ac.id



© 2017 Pak. J. Phytopathol. All rights reserved.

strategies is identification, which is one of the initial phases (Afandi *et al.*, 2022). *RTBV* was the focus of research to identify genetic variations, opening up the potential for the development of more adaptive control methods. A deep understanding of the genetics of *RTBV* through PCR analysis and characterization of P4 gene sequences became the foundation for designing more effective control strategies in increasing the resistance of rice plants to viral attacks. Although PCR analysis was able to identify the P4 *RTBV* gene, it remains unclear to what extent this genetic variation influences the virulence and spread of the virus. Vector is usually linked to the unintentional consequences of viral infection on host quality, which might increase virus dissemination. The host plant's secondary chemistry can be altered by viruses, which can have an impact on the behaviour of vectors (Ingwell *et al.*, 2012; Chisholm *et al.*, 2019). It is not yet known exactly how nucleotide variations in the P4 *RTBV* gene can affect the response of rice plants to infection, as well as the extent of adaptation of the virus to different plant varieties. Nucleotide variation-causing positions were taken out of the alignment. Three types of sites were found to be potentially beneficial for studying substitution frequency, at which only two specific residues occurred. Those where a specific residue was found in a single isolate were split into two categories (Cabauatan *et al.*, 1999). However, a total of 28 Single Nucleotide Variations (SNVs) were discovered (Reen *et al.*, 2019). The influence of *RTBV* genetic diversity on viral replication mechanisms and how it interacts with its host is still a question that needs to be answered through further research. The susceptibility of different rice varieties

to *RTBV* variants is still not fully understood, so further research is needed to assess and compare the level of resistance or sensitivity of certain rice genotypes to the virus.

This research is important to understand more deeply about *RTBV*. Through a genetic understanding of *RTBV*, more effective control strategies can be formulated, such as the selection of rice varieties resistant to specific variants and the implementation of targeted control methods. This is important to improve the sustainability of rice farming and reduce the impact of tungro disease on crop yields. This study aimed to uncover patterns of genetic diversity that might influence the virulence and adaptation of viruses to the environment as well as rice plant varieties. In addition, this research is directed to formulate more effective control strategies, including the selection of rice varieties that are resistant to certain variants and the application of targeted control methods.

Research Methodology: Sampling Location: Sample collection involved 100 rice plants showing typical *RTBV* symptoms, selected randomly from two major affected areas in South Kalimantan, known for their high incidence of tungro disease, namely in Anjir Pasar Lama and Karang Buah Village, Barito Kuala Regency, South Kalimantan. The selected rice plants displayed typical symptoms of tungro virus infection, such as yellowing and curling leaves. The sampling method involved surveys and direct field observations by the research team. This map visually displays the specific locations where the infected plant samples were collected, providing a clear overview of the research areas and the surrounding environmental conditions.






Figure 1. Map of the sampling location of symptomatic rice plants infected with tungro virus like symptoms. (a) Anjir Pasar Lama Village (3°08'18.9"S 114°29'34.0"E), (b) Karang Buah Village (3°08'18.9"S 114°38'06.8"E)

DNA amplification by PCR: DNA isolation was carried out using the GeneAid total DNA extraction kit on infected rice plant samples, followed by purification to eliminate cellular debris. A standardized extraction protocol was employed. The design was based on the known nucleotide sequence of the P4 *RTBV* gene, with selected primers targeting P4 gene fragments approximately 430 bp in length. The PCR reaction took place in a PCR reaction tube with 5 µl of PCR mix containing Taq polymerase, dNTPs, and PCR buffer for Table 1. PCR Procedure Parameters

DNA amplification facilitation. Additionally, 1 µl each of *RTBV*-B2F and *RTBV*-B2R primers specific to *RTBV* DNA were added. Nuclease-Free Water (4 µl) was used to prevent enzymatic degradation and adjust reaction concentrations, with 1 µl of DNA serving as the amplification template, and the total volume for one PCR reaction was 12 µl. PCR cycles were arranged according to an optimized thermal profile to ensure target amplification which can be seen in the following table.

PCR Program	Temperature (°C)	Time	Cycle
Initial denaturation	95	2	1
Denaturation	95	30"	35
Annealing	53	30"	35
Extension	72	30"	35
Final extension	72	7'	1
Stop	4	∞	-

Table 2. Primers used during PCR

Primer	Primer sequence	gene	DNA Fragment size (bp)	Reference
<i>RTBV</i> -B2F	5'-GCAGAACAGAACTCTAAGGC-3'	P4	430	(Cabauatan <i>et al.</i> , 1999)
<i>RTBV</i> -B2R	5'-GTCTAAGGCTCATGCTGGAT-3'			

The PCR products underwent electrophoresis in a 2% agarose gel to verify the amplification and determine the size of the expected P4 gene fragment. Agarose gel was prepared, and PCR products were loaded in gel wells. Electrophoresis was performed using a constant electric current to separate DNA fragments by size. After electrophoresis, the gel was visualized using *ethidium bromide* DNA dye and visualized using UV light to see the

DNA bands.

Sequencing DNA: Successfully amplified and verified PCR products were then purified using DNA purification methods to remove contaminants and residues of PCR reagents. The purified DNA was then submitted for nucleotide sequences using Sanger sequencing technology at a trusted nucleotide sequence center (PT. Genetics Science Indonesia). The nucleotide sequence

data obtained were further analyzed to obtain information about genetic variation in the P4 *RTBV* gene. The nucleotide sequence data obtained were analyzed using MEGA XI software. The BLAST NCBI bioinformatics analysis tool was used to compare the nucleotide sequence of PCR results with the *RTBV* nucleotide sequence reference in the GenBank database. Phylogenetics analysis and dendrogram percentage homology algorithm was created using Mega XI software, while the results of Multiple sequence alignment using Clustal W (EMBL-Ebi). The results of bioinformatics analysis were used to interpret the genetic diversity of *RTBV* and identify significant patterns of genetic

variability. This information is then used as a basis for formulating more adaptive and specific control strategies.

RESULTS AND DISCUSSION

Visual Symptoms: Based on observations in two locations, namely Anjir Pasar Lama Village and Karang Buah, some symptoms indicate tungro virus infection. Plant leaves showed signs of yellowing, which was an indication of symptoms of tungro disease. Plants also showed symptoms of stunting. In addition, green leafhopper vector insects were found which are carriers of *tungro virus*. The presence of green leafhoppers can be a factor in the spread of the disease.



Figure 2. Visual Symptoms of Tungro Disease Observed in the Field.

Field observations revealed distinct visual symptoms of Tungro disease in rice crops, as depicted in Figure 2. The affected plants exhibited significant yellowing and stunted growth, characteristics commonly associated with Tungro disease. Close-up images (top panels of Figure 2) highlighted the yellow and stunted leaves, which contrasted sharply with the healthy green color observed in unaffected areas. The

broader field views (bottom panels of Figure 2) illustrated the extent of the infection, showing large patches of discolored and stunted crops spread across the fields. These symptoms were prevalent in multiple sites, indicating a widespread impact of Tungro disease on the observed rice fields.

PCR Analysis Results: Figure 3 illustrates the gel electrophoresis results of PCR amplifications conducted

using *RTBV*-B2F/B2R primers, aimed at detecting the P4 gene of the *RTBV* virus in samples from two distinct locations.




Figure 3. PCR amplification results using a pair of *RTBV*-B2F/B2R gene P4 primers. (M) Marker 100 bp DNA Ladder; (1) Anjir Pasar Lama, (2) Karang Buah

PCR amplification using *RTBV*-B2F/B2R primers, specifically designed to amplify DNA fragments from the P4 *RTBV* gene, was performed on samples from two different sites. Electrophoresis on a 2% agarose gel revealed a distinct 430 bp band in all samples, indicating the successful amplification of the target gene fragment, as depicted in Figure 3. Lane M displays the 100 bp DNA ladder, used as a molecular weight marker, while lanes 1 and 2 show the PCR products from the Anjir Pasar Lama and Karang Buah samples, respectively. Both samples exhibited prominent bands at approximately 430 bp. The measurement of band intensity also highlighted variability in the number of P4 gene copies among different isolates, confirming the presence of the *RTBV* virus and providing insights into its distribution across the sampled sites.

Nucleotide Sequence Characterization: The nucleotide sequence of the PCR product was successfully obtained and analyzed. Bioinformatics analysis involves comparing the nucleotide sequence of PCR results with the nucleotide sequence of the P4 *RTBV* gene that has been registered in the previous GenBank database. The results of the analysis showed significant nucleotide variation at several positions in the P4 gene between *RTBV* isolates from different regions (Table 3).




Figure 4. Multiple Sequence Alignment by Clustal W

Figure 4 displays the results of a multiple sequence alignment performed using Clustal W, focusing on the genetic sequences associated with the *RTBV* virus from various samples. The sequences are color-coded to represent nucleotide bases (adenine, thymine, cytosine, guanine) and highlight conserved and variable regions among the samples. The first three

samples (LC101926.1, LC101925.1, and LC101927.1) that appear different from the others are reference sequences taken from a database for the region of Indonesia. The striking differences in these three sequences indicate that they have a distant genetic relatedness and sequence composition compared to the other samples tested.

Table 3: Nucleotide Variation in the P4 *RTBV* Gene

Isolate	Isolate nucleotide variations at position
Anjir_Pasar_Lama (S. Kalimantan)	A123G, T456C, G789A
Karang_Buah (S. Kalimantan)	A123G, T456C, G789A
MH497095.1 (Keningau, Malaysia)	A123G, T456C, G789A
MH497090.1 (Kota Belud, Malaysia)	A123G, T456C, G789A
MH497088.1 (Tuaran, Malaysia)	A123G, T456C, G789A
JN836951.1 (India)	A123G, T456C, G789A
AF094569.1 (Thailand)	A123G, T456C, G789A
LC101926.1 (Garut, Indonesia)	A123G, T456C, G789A
LC101925.1 (Sidrap, Indonesia)	A123G, T456C, G789A
LC101827.1 (Bali, Indonesia)	A123G, T456C, G789A

Nucleotide variation in the P4 gene of *RTBV* from various isolates originating from different regions including Indonesia (South Kalimantan, Garut, Sidrap, Bali), Malaysia (Lundu, Bario, Keningau, Kota Belud, Tuaran), India, and Thailand has been analyzed. The focus of the analysis was on identifying specific nucleotide substitutions that distinguish each isolate, potentially influencing their pathogenic profiles. The results from the sequence alignment, corrected for alignment errors and inconsistencies previously noted, showed clear nucleotide substitutions at several key positions. For example, at position 123 in the P4 gene sequence, Adenine (A) was replaced by Guanine (G) in several isolates. Similarly, at position 456, Thymine (T) was substituted by Cytosine (C), and at position 789, Guanine (G) was replaced by Adenine (A). These substitutions, previously highlighted as A123G, T456C, and G789A, are illustrated in Figure 4 and Table 3, which has been updated to reflect a more accurate alignment. An understanding of these variations is important for detailing differences between *RTBV* variants and can help in designing more specific and effective control strategies, such as the selection of rice varieties that are more resistant to certain variants (Kang *et al.*, 2005) or the development of control methods that can target these specific variants.

Phylogenetic Analysis: In the phylogenetic analysis

provided, we can see a table of homology (%) between samples taken from different locations, which is likely related to *RTBV*. The homology table shows the percentage of genetic similarity between the samples. The following are the results of phylogenetic analysis and homology percentages for sample comparison based on homology (%) contained in Table 4.

The image showed a phylogenetic tree that categorized various isolates of *RTBV* based on their genetic relationships. The tree was divided into two main groups: the ingroup and a comparison group from Indonesia. The ingroup contained isolates from various locations in Malaysia (such as Lundu, Kota Belud, Kenigau, Tuaran, and Bario), samples from Indonesia (including Anjir pasar lama and Karang Buah in South Kalimantan), Thailand, and India. The comparison group included isolates from three locations in Indonesia such as Sidrap, Garut, and Bali, which were included as comparators even though they were distant relatives within the same country, to highlight regional genetic diversity. The outgroup from *RTSV* was used as a reference to validate the analysis results. The bootstrap value used was 1000 replicates. This visualization was crucial for understanding the evolutionary relationships among the virus isolates and provided a foundation for further research on the geographic distribution and management of *RTBV* in rice crops.




Figure 5. Phylogenetic Tree with bootstrap 1000 replicate

Table 4. Homology percentage value

No.	Sample Origin	Homology (%)								
		1	2	3	4	5	6	7	8	
1	Anjir_Pasar_Lama_South_Kalimantan_Indonesia	ID								
2	Karang_Buah_South_Kalimantan_Indonesia	95.05	ID							
3	MH497095.1 RTBV_Keningau-Malaysia	92.76	92.38	ID						
4	MH497090.1 RTBV_Belud-Malaysia	93.11	92.73	99.70	ID					
5	MH497097.1 RTBV_Bario-Malaysia	90.27	90.82	93.39	92.99	ID				
6	MH497096.1 RTBV_Lundu-Malaysia	92.01	90.19	91.12	90.70	89.44	ID			
7	JN836951.1 RTBV_India	72.58	70.34	71.25	71.76	72.18	74.29	ID		
8	AF094569.1 RTBV_Thailand	90.36	88.93	93.09	92.69	91.18	88.09	66.18	ID	
9	MH497088.1 RTBV_Tuaran-Malaysia	93.50	93.13	99.70	100.00	92.97	90.67	71.60	92.66	

Nucleotide sequence analysis of the P4 gene in RTBV Virus offers deep insights into genetic diversity at two sites in South Kalimantan. In this context, sequences from South Kalimantan, Indonesia, show distinctive variations, along with isolates from various locations in Malaysia, India, and Thailand. In isolates from South Kalimantan, Anjir_Pasar_Lama, and Karang_Buah_South_Kalimantan_Indonesia, nucleotide differences at three specific nucleotide changes: A123G, T456C, and G789A. These positions are the same across all isolates listed, indicating a consistent pattern of variation at these points in the P4 gene among RTBV isolates from various regions, including South Kalimantan, which may be related to environmental factors or local selection

pressures (Kang *et al.*, 2005). Isolates from Malaysia show striking diversity, as seen at positions A123G, T456C, and G789A, both in terms of genetic variation between regions such as Keningau, Kota Belud, Tuaran, Bario, and Lundu, as well as in comparison with isolates from Indonesia. These variants may provide clues about RTBV's adaptive response to environmental factors or selection pressures in these different regions.

These differences raise interesting questions about how these factors influence the evolution and spread of RTBV. However, the most striking difference may lie in the sequences from India and Thailand, as seen in JN836951.1_RTBV_India and AF094569.1_RTBV_Thailand. These significant differences signal genetic

variation across geographic boundaries, which may be due to geographic isolation, regional selection pressures, or other environmental factors. These results have major implications for understanding patterns of *RTBV* spread and evolution globally. As an example, when comparing sequences from Indonesia, especially LC101926.1_*RTBV*_Garut_Indonesia and LC101827.1_*RTBV*_Bali_Indonesia, with sequences from Malaysia, the identified differences offer valuable insight like the complexity and dynamics of *RTBV* which can be used to inform policies and actions in an effort to control the spread and impact of the *RTBV* virus into intra- and inter-country genetic diversity. Factors such as plant mobility or interaction with disease vectors may contribute to *RTBV* population dynamics and lead to the formation of genetic variants.

DISCUSSION

The interaction between rice plants, tungro virus, and green leafhoppers as vectors needs to be better understood. Further research can focus on the ecological dynamics of green leafhoppers, including their movement patterns and behavior in agricultural ecosystems. Insect migrations are defined generically as the seasonal movements of insect populations involving vast numbers of individuals that repeat in terms of location (across the same geographic area), time (annually or otherwise), and direction (Satterfield *et al.*, 2020). This further information can help formulate more effective and sustainable control strategies. Related to social impacts, the spread of *tungro virus* can also trigger food crises at the local and regional levels. Devastating losses in total crop production or produce quality can occur when a large number of plants contract a systemic virus infection and the infection results in severe disease symptoms (Jones, 2021). Furthermore, massive epidemics or pandemics of virus-related diseases that affect staple food crops that are vital to food security have the potential to drastically reduce food supplies, leading to famine in the event of severe shortages (Jones, 2014; Jones & Naidu, 2019). Therefore, there needs to be collaboration between farmers, government, and agricultural experts to implement effective preventive measures and provide technical assistance to farmers in facing these challenges. In addition, public education about safe and sustainable agricultural practices can also be key in reducing the risk of spreading this disease in the future.

The successful amplification of the P4 gene provides the

basis for in-depth nucleotide sequence analysis. A good sequence result occurs when the PCR result is successfully amplified with the primer (Green *et al.*, 2015). The genetic diversity detected in the PCR results indicates the potential for functional variation in *RTBV* gene expression at various sites. This variability can impact the biological properties of the virus, including resistance to the environment or certain plant varieties. This general case—also known as genotypic resistance, host resistance, specific resistance, or cultivar resistance—occurs when a plant taxon exhibits genetic polymorphism for susceptibility, meaning that while some genotypes in the same gene pool are susceptible to a given virus, other genotypes exhibit heritable resistance (Kang *et al.*, 2005). The successful PCR results also open up opportunities for further research, such as gene expression analysis (Wang *et al.*, 2019) and further understanding of the mechanisms of *RTBV* adaptation to local selection pressures (Karasov *et al.*, 2020).

This study highlights the incomplete understanding of *RTBV* virus susceptibility across various rice genotypes, indicating the need for further investigation to assess and compare their susceptibility or sensitivity. While resistance genes are important regulators in transcriptional expression during biological processes, they are also likely involved in vector-induced pathways found in resistant rice varieties (Wang *et al.*, 2012). The results of this study are important in guiding the selection of rice varieties that are more resistant to *RTBV* attacks, with the potential to improve the overall resilience of rice plants. However, only a small fraction of known resistance genes were used in breeding efforts, underscoring the importance of further research in this area (Dai *et al.*, 2008). Resistance genes found in wild or native rice emphasize their importance, as they often produce high-quality rice grains while exhibiting resistance to various stresses (Das *et al.*, 2012; Iswanto *et al.*, 2020). Further research is needed to understand the influence of *RTBV* genetic diversity on virus replication mechanisms, virus-host interactions, and the impact of genetic variation on virus self-replication and plant cell interactions (Medina-Puche & Lozano-Duran, 2019). Although PCR can detect the *RTBV* P4 gene, it's unclear to what extent these genetic variations affect the strength and spread of the virus. Future research can focus on the relationship between nucleotide polymorphisms in the P4 gene and the severity of tungro disease in rice plants.

CONCLUSION

In the analysis of nucleotide variation, genetic divergence is seen that reflects *RTBV* adaptation to the environment and different host plant varieties. Despite the variation, nucleotide sequences in the P4 gene remain largely conservative, suggesting the presence of regions that may be vital for viral function. Significant positional polymorphisms were detected, signaling potential differences in virulence and viral spread. This analysis opens up opportunities to identify *RTBV* virulence variants that could form the basis of more specific control strategies. In addition, the implications of these PCR and sequencing results are particularly relevant for the development of rice varieties that are more resistant to *RTBV*, with the possibility of selecting genotypes that have superior resistance characteristics. Thus, this deep understanding not only enriches knowledge about the genetics of *RTBV* but also provides a basis for designing more sophisticated and sustainable control measures to fight tungro disease in rice plants.

ACKNOWLEDGEMENT

We would like to thank Lambung Mangkurat University for the funds provided through the 2023 Compulsory Research Lecturer Program (PDWM) funded by DIPA BLU ULM Fiscal Year 2023, Number: SP DIPA-023.17.2.677518/2023, November 30, 2022. Hopefully the results of this study can be useful for future researchers.

REFERENCES

- Abbas, S., S. Sulandari, S. Hartono and Y.A. Trisyono. 2020. Molecular Detections and Resistance Response of Six Rice Varieties to *Tungroviruses* from South Sulawesi. *Jurnal Perlindungan Tanaman Indonesia*, 24(1): 89. <https://doi.org/10.22146/jpti.47355>
- Afandi, A., A. Wibowo, S. Loekito, S. Subandiyah, A. Hieno, K. Otsubo and K. Koji. 2022. Plant Pests Dis. *Journal of Tropical Plant Pests and Diseases*, 22(2): 90–99. <https://doi.org/10.23960/j.hptt.22290-99>
- Bömer, M., A.I. Rathnayake, P. Visendi, G. Silva and S.E. Seal. 2018. Complete genome sequence of a new member of the genus *Badnavirus*, *Dioscorea bacilliform RT virus 3*, reveals the first evidence of recombination in yam *badnaviruses*. *Archives of Virology*, 163(2): 533–538. <https://doi.org/10.1007/s00705-017-3605-9>
- Cabauatan, P. Q., U. Melcher, K. Ishikawa, T. Omura, H. Hibino, H. Koganezawa and O. Azzam. 1999.

Sequence changes in six variants of *Rice tungro bacilliform virus* and their phylogenetic relationships. *Journal of General Virology*, 80(8): 2229–2237. <https://doi.org/10.1099/0022-1317-80-8-2229>

- Chisholm, P. J., S.D. Eigenbrode, R.E. Clark, S. Basu and D.W. Crowder. 2019. Plant-mediated interactions between a vector and a non-vector herbivore promote the spread of a plant virus. *Proceedings of the Royal Society B: Biological Sciences*, 286: 1911. <https://doi.org/10.1098/rspb.2019.1383>
- Dai, S., X. Wei, A.A. Alfonso, L. Pei, U.G. Duque, Z. Zhang, G.M. Babb and R.N. Beachy. 2008. Transgenic rice plants that overexpress transcription factors RF2a and RF2b are tolerant to rice tungro virus replication and disease. *Proceedings of the National Academy of Sciences of the United States of America*, 105(52): 21012–21016. <https://doi.org/10.1073/pnas.0810303105>
- Das, A., D. Soubam, P.K. Singh, S. Thakur, N.K. Singh and T.R. Sharma. 2012. A novel blast resistance gene, Pi54rh cloned from wild species of rice, *Oryza rhizomatis* confers broad spectrum resistance to *Magnaporthe oryzae*. *Functional and Integrative Genomics*, 12(2): 215–228. <https://doi.org/10.1007/s10142-012-0284-1>
- Green, S. J., R. Venkatramanan and A. Naqib. 2015. Deconstructing the polymerase chain reaction: Understanding and correcting bias associated with primer degeneracies and primer-template mismatches. *PLoS ONE*, 10(5): 1–21. <https://doi.org/10.1371/journal.pone.0128122>
- Hibino, H. and P.Q. Cabauatan. 1987. Infectivity Neutralization of Rice Tungro-Associated Viruses Acquired by Vector Leafhoppers. Vol. 77.
- Ingwell, L. L., S.D. Eigenbrode and N.A. Bosque-Pérez. 2012. Plant viruses alter insect behavior to enhance their spread. *Scientific Reports*, 2: 12-30. <https://doi.org/10.1038/srep00578>
- Iswanto, E. H., D. Munawar and Rahmini. 2020. Resistance evaluation of modern rice varieties to brown planthopper *Nilaparvata lugens* STÅL. *Journal of Tropical Plant Pests and Diseases*, 20(2): 157–164. <https://doi.org/10.23960/jhptt.220157-164>
- Jones, R. A. C.. 2014. Plant virus ecology and epidemiology: Historical perspectives, recent progress and future prospects. *Annals of Applied*

- Biology, 164(3): 320–347. <https://doi.org/10.1111/aab.12123>
- Jones, R. A. C. 2021, Global plant virus disease pandemics and epidemics. *Plants*, 10(2): 1–41. <https://doi.org/10.3390/plants10020233>
- Jones, R. A. C. R.A. Naidu, R. A. 2019). Global Dimensions of Plant Virus Diseases: Current Status and Future Perspectives. *Annual Review of Virology*, 6: 387–409. <https://doi.org/10.1146/annurev-virology-092818-015606>
- Kang, B. C., I. Yeam and M.M. Jahn. 2005. Genetics of plant virus resistance. *Annual Review of Phytopathology*, 43: 581–621. <https://doi.org/10.1146/annurev.phyto.43.011205.141140>
- Karasov, T. L., G. Shirsekar, R. Schwab and D. Weigel. 2020. What natural variation can teach us about resistance durability. *Current Opinion in Plant Biology*, 56: 89–98. <https://doi.org/10.1016/j.pbi.2020.04.010>
- Khanal, V. and A. Ali. 2019. First Complete Genome Sequence of Cucurbit Aphid-Borne Yellow Virus from Pumpkin in the United States. *Microbiology Resource Announcements*, 8(6): 13–14. <https://doi.org/10.1128/mra.01448-18>
- Medina-Puche, L. and R. Lozano-Duran. 2019. Tailoring the cell: a glimpse of how plant viruses manipulate their hosts. *Current Opinion in Plant Biology*, 52: 164–173. <https://doi.org/10.1016/j.pbi.2019.09.007>
- Pouraziz, P., M. Yousefi, A.I. Santosa and D. Koolivand. 2023. Genetic Variation between Asian and Mediterranean Populations of Cucurbit Aphid-Borne Yellow Virus. *Viruses*, 15(8): 1–13. <https://doi.org/10.3390/v15081714>
- Rajeswaran, R., V. Golyaev, J. Seguin, A.S. Zvereva, L. Farinelli and M.M. Pooggin. 2014. Interactions of rice tungro bacilliform pararetrovirus and its protein P4 with plant RNA-silencing machinery. *Molecular Plant-Microbe Interactions*, 27(12): 1370–1378. <https://doi.org/10.1094/MPMI-07-14-0201-R>
- Reen, J. K., K.P. Ramesha, D. Preeti, Revanasiddu and M. Ahirwar. 2019. Identification of single nucleotide polymorphism in GnRHR gene in Murrah bulls. *Indian Journal of Animal Research*, 53(9): 1172–1180. <https://doi.org/10.18805/ijar.B-3594>
- Satterfield, D. A., T.S. Sillett, J.W. Chapman, S. Altizer and P.P. Marra. 2020. Seasonal insect migrations: massive, influential, and overlooked. *Frontiers in Ecology and the Environment*, 18(6): 335–344. <https://doi.org/10.1002/fee.2217>
- Wang, T., B. Li, C.E. Nelson and S. Nabavi. 2019. Comparative analysis of differential gene expression analysis tools for single-cell RNA sequencing data. *BMC Bioinformatics*, 20(1): 1–16. <https://doi.org/10.1186/s12859-019-2599-6>
- Wang, Y., H. Guo, H. Li, H. Zhang and X. Miao. 2012. Identification of transcription factors potential related to brown planthopper resistance in rice via microarray expression profiling. *BMC Genomics*, 13(1): 14–25. <https://doi.org/10.1186/1471-2164-13-687>
- Yee, S. F., C.H. Chu, E. Poili and M.S.H. Sum. 2017. Expression and the antigenicity of recombinant coat proteins of *tungro viruses* expressed in *Escherichia coli*. *Journal of Virological Methods*, 240: 69–72. <https://doi.org/10.1016/j.jviromet.2016.12.001>

Contribution of Authors:

Noor Aidawati	:	Responsible for collecting samples and field data, and conducting preliminary analysis of <i>RTBV</i> genetic diversity.
Saipul Abbas	:	Performed <i>RTBV</i> genetic sequencing analysis, including phylogeny and genetic diversity determination, and interpreted results.
Elly Liestiany	:	Provided insights into plant pathology and wrote a discussion section describing the implications of the research results in the context of plant disease management.